
## NÜROO POCKET VALUE ANALYSIS



Hospital Collection

www.brownmed.com

## REFERENCES

Feldman R, Rosenthal Z, Eidelman AI. Maternal-preterm skin-to-skin contact enhances child physiologic organization and cognitive control across the first 10 years of life. Biol Psychiatry. 2014 Jan 1;75(1):56–64.

Conde-Agudelo A, Belizán JM, Diaz-Rossello J. Kangaroo mother care to reduce morbidity and mortality in low birthweight infants. In: The Cochrane Collaboration, editor. Cochrane Database of Systematic Reviews [Internet]. Chichester, UK: John Wiley & Sons, Ltd; 2011 [cited 2016 Apr 12]. Available from: http://doi.wiley.com/10.1002/14651858.CD002771.pub2

Samra NM, Taweel AE, Cadwell K. Effect of intermittent kangaroo mother care on weight gain of low birth weight neonates with delayed weight gain. J Perinat Educ. 2013;22(4):194–200.

Johnston C, Campbell-Yeo M, Fernandes A, Inglis D, Streiner D, Zee R. Skin-to-skin care for procedural pain in neonates. Cochrane Database Syst Rev. 2014;1:CD008435.

Length of Stay - Reducing Length of Stay - NHS Institute for Innovation and Improvement [Internet]. [cited 2016 Apr 12]. Available from: http://www.institute.nhs.uk/quality\_and\_service\_improvement\_ tools/quality\_and\_service\_imporvement\_tools/length\_of\_stay/

McClead RE MD. Length of Stay Reduction: A Quality Improvement Project for Neonatal Abstinence Syndrome [Internet]. Available from: http://www.astho.org/Programs/Prevention/Injury-and-Violence-Prevention/\_Materials/Neonatal\_Abstinence\_Syndrome\_MCCLEAD/